


Geosur 2007

Un Congreso Internacional sobre Geología y Geofísica
del Hemisferio Sur

Santiago de Chile
18 al 20 de Noviembre 2007

Libro de Resúmenes

Editores : A. Demant, F. Hervé, M. Menichetti and A. Tassone



Organizan :

Departamento de Geología, FCFM, Universidad de Chile
Proyecto ARTG-04, "Conecciones Antártica - Patagonia", PBCT-CONICYT
Departamento de Ciencias Ecológicas, Pontificia Universidad Católica de Chile

Auspician:



Antofagasta
Minerals



Empresa Nacional
del Petróleo

Una contribución a :

Año Internacional del Planeta Tierra
Año Polar Internacional

Famatinian cordierite-bearing magmatism on the proto-Andean margin of Gondwana

Alasino P. ^{*1}, Dahlquist J. ², Rapela C. ³, Galindo C. ⁴, Casquet C. ⁴, Pankhurst R. ⁵, Saavedra J. ⁶ and Baldo E. ².

⁽¹⁾ CRILAR-CONICET, Entre Ríos y Mendoza, 5301, Anillaco, La Rioja, Argentina

⁽²⁾ UNC-CONICET, Av. Vélez Sársfield 1611, Pabellón Geol., X5016CGA-Argentina

⁽³⁾ CIG-UNLP, Calle 1 N° 644, B1900TAC, La Plata, Argentina.

⁽⁴⁾ Depto. de Petrología y Geoquímica, UCM, 28040 Madrid, España

⁽⁵⁾ BGS, Kingsley Dunham Centre, Keyworth, Nottingham NG12 5GG, UK

⁽⁶⁾ IRNASA-CSIC, Cordel de Merinas 40-52, 257 37008, Salamanca, España

* Presenting author email: palasino@crilar-conicet.com.ar

Petrological and geochemical data from the Cerro Toro unit and Mazán granite, in Sierra de Famatina and Sierra de Mazán, respectively, as well as the Tuani Granite in the Sierra de Chepes (Dahlquist *et al.*, 2005), indicate that the Famatinian peraluminous magmatism in the Sierras Pampeanas defines trends from intermediate (60.7-69.9%) to felsic (71.3-73.6%) compositions, with mineral associations formed by Pl, Mc, Qtz, Bt, Ms, Crd, \pm Gtr and Zrn, Mon, fluor-Ap, Opq (mainly ilmenite and/or titanomagnetite). The granites show a distinctive high Aluminium Saturation Index (1.39-2.07), that increases towards the most mafic varieties defined by cordierite-bearing granitoids with high $\text{FeO}_t + \text{MgO} + \text{TiO}_2$ (5.7-11.3%) and intermediate $\text{Na}_2\text{O} + \text{K}_2\text{O}$ (5.2-6.7%) contents. All granitic suites define similar magmatic trends in the high peraluminous field of the A-B multicationic diagram, indicating that they crystallized in medium- to low- P conditions ($A = 60$ -180 and $B = 100$ -190), similar to those observed in other cordierite bearing S-type granites (*e.g.*, Cooma granodiorite in the Lachlan Fold Belt, SW Australia). Moreover, source discrimination diagrams show that the trends of the cordierite-bearing suites are projected in the greywacke field, with $\text{Rb/Ba} = 0.16$ -0.56, $\text{Rb/Sr} = 0.70$ -2.83 and $\text{CaO/Na}_2\text{O} = 0.34$ -0.69. Euhedral crystals of magmatic cordierite [$\text{Mg}/(\text{Mg} + \text{Fe} + \text{Mn}) = 0.59 \pm 0.04$; $\text{Na} + \text{K} = 0.05 \pm 0.01$, $n = 23$], devoid of mineral inclusions, are typical of all these granites, accompanied by plagioclase ($\text{An} = 29.7 \pm 4.8$, $n = 23$), alkali feldspar ($\text{Or} = 84.9 \pm 7.1$, $n = 6$), biotite [$\text{Fe}/(\text{Fe} + \text{Mg}) = 0.53 \pm 0.06$, $n = 31$] and garnet ($\text{Alm} = 72.2 \pm 1.3$, $\text{Prp} = 12.2 \pm 1.6$ and $\text{Spss} = 12.2 \pm 2.7$, $n = 10$).

This peraluminous magmatism is the result of regional anatexis of a plagioclase-rich, immature continental platform sedimentary source, during a time-restricted episode (484-479 Ma, U/Pb SHRIMP, Pankhurst *et al.*, 2000), and with a high geothermal gradient.

References

- Dahlquist J.A., Rapela C.W., Baldo E., 2005. Journal of South American Earth Science **20**, 231-251.
- Pankhurst R., Rapela C., Fanning C., 2000. Transactions of the Royal Society of Edinburgh: Earth Sciences **91**, 151-168.